

PRECAUTIONS

Store in a well-ventilated area, away from direct sunlight, food, and feeds. Keep away from reach of children. Wear protective clothing while handling the product

BABALA: Itago ang produkto sa angkop na lugar, malayo sa init ng araw. Ihiwalay sa pagkain ng tao o hayop. Magsuot ng proteksyong pananamit habang ginagamit ang produkto

IT IS A VIOLATION OF FPA RULES & REGULATIONS TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABEL.

SYMPTOMS OF POISONING:

No case of human poisoning is on record.

NOTE TO PHYSICIAN:

No special antidote. Treat symptomatically and supportively.

FIRST AID:

In case of contact with skin, rinse the area thoroughly with soap and water. In case of contact with eyes, rinse thoroughly in running water for 15 minutes and immediately consult a doctor. After spraying, thoroughly wash hands and all other body parts which have been in contact with the material. If medical advice is needed, have product container or label at hand

Manufactured by:

STOCKTON (ISRAEL) LTD.

17 Ha Mefalsim St. P.O.B. 3517. Petach Tikva 4951447, Israel

Distributed by:

 **AGWAY CHEMICALS CORPORATION**

Address: GFR Building, Lot 9B, JP Laurel Avenue, Davao City, 8000
Telephone Nos.: (082) 226-3668; (082) 300-9515
Website: <https://agwaychemicals.com.ph>

Formulation Date: (dd/mm/yyyy)

Batch Number:


bio-ag technologies

Member:



GROUP F7 FUNGICIDE
GROUP G1 FUNGICIDE

stk REGEV™

FIRST HYBRID FUNGICIDE

For the control of:
Black Sigatoka in Banana,
Panicle / Blossom Blight, Anthracnose
and Stem-End Rot in Mango
Sheath Blight & Blast in Rice

Composition

Active Ingredients:

Tea Tree Oil **400 g/L**
(an extract of *Melaleuca Alternifolia*)

Difenoconazole **200 g/L**

Inert Ingredients **400 g/L**

Total 1000 g/L

Net Content: 1000 mL

Registered with the Fertilizer and Pesticide Authority Pursuant to P.D. 1144
FPA Registration No.: **527-480-4399**

Imported by:

STOCKTON - AGRIMOR PHILIPPINES INCORPORATED

Unit 28, D3G Y10 Bldg., Mamay Road, Angliongto Ave., Davao City, 8000
Tel. No. (082) 327 3830

KEEP OUT OF REACH OF CHILDREN

BEFORE APPLYING THE PRODUCT, READ AND UNDERSTAND THIS LABEL

GENERAL INFORMATION:

STK REGEV is the first hybrid fungicide combining a plant extract and a systemic fungicide in a single formulation. It is ideal for Integrated Pest Management (IPM) programs.

RECOMMENDATION:

Spray during the early hours of the day. Avoid spraying in temperatures above 35° C. No phytotoxicity has been observed when product is applied according to the recommended selectivity testing procedure.

| Crop | Disease(s) | Dose rate (mL/ha) | Spray volume (L/ha) |
|--------|---------------------------------------------------------|----------------------------------|-----------------------------|
| Banana | Black Sigatoka | 300 - 500 | 25 - 30 Aerial Application |
| | | | 80 - 100 Ground Application |
| Mango | Anthracnose Panicle / Blossom blight Stem-End Rot | 1000 - 1250 100 - 125 mL/200L | 2000 200L Power Sprayer |
| | | | |
| Rice | Sheath Blight Rice Blast | 250 - 500 20-40 mL/16L | 200 16L Sprayer |

PRE-HARVEST INTERVAL:

Banana: Zero (0) days | Mango: Fourteen (14) days | Rice: Seven (7) days

RE-ENTRY PERIOD: As soon as foliage dries up

COMPATIBILITY: STK REGEV is compatible with commonly used fungicides. It is recommended to carry out tests before mixing with any other product. The manufacturer and distributor should be consulted before mixing with other products and adjuvants.

NOTICE: No warranty of any kind, expressed or implied, is made concerning the use of this product. User assumes all risks and liabilities resulting from handling, use or application

STORAGE AND DISPOSAL: Store in closed, original container in a cool, dry, well-ventilated area. **DO NOT** store for prolonged periods in direct sunlight or damp area. Triple rinse the empty containers before disposal. Add rinsate to spray tank and pour on non-crop area. **DO NOT** dispose undiluted chemicals on-site. Break crush or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. **DO NOT** burn empty containers. Dispose of containers, in accordance with local government regulations.

IN CASE OF EMERGENCY, PLEASE CONTACT

National Poison Control and Information Service
PGH Manila
Tel. (02) 521-8451 Loc. 2311
(02) 524-1078

Vicente Sotto Medical Center
B. Rodriguez St. Cebu City
Tel. (032) 233-9891 loc. 348

Southern Philippines Medical Center
Bajada, Davao City
Tel. (082) 227-2731



STK REGEV 1L label height: 140mm width 280mm